

WHAT IS CLAIMED IS:

Claim 1. A method for the imaging of a particular volume of plant or animal tissue, wherein the plant or animal tissue contains at least one photo-active molecular agent. the method comprising the steps of:

- 5
- (a) treating the particular volume of the plant or animal tissue with light sufficient to promote a simultaneous two-photon excitation of the photo-active molecular agent contained in the particular volume of the plant or animal tissue;
 - (b) photo-activating at least one of the at least one photo-active molecular agent in the particular volume of the plant or animal tissue, thereby producing at least one photo-activated molecular agent, wherein the at least one photo-activated molecular agent emits energy;
 - (c) detecting the energy emitted by the at least one photo-activated molecular agent; and
 - (d) producing a detected energy signal which is characteristic of the particular volume of plant or animal tissue.

Claim 2. The method of Claim 1 wherein the light sufficient to promote a simultaneous two-photon excitation of the at least one photo-active molecular agent is laser light.

Claim 3. The method of Claim 1 wherein the light sufficient to promote a simultaneous two-photon excitation of the photo-active molecular agent is a focused beam of light.

Claim 4. The method of Claim 4 wherein the focused beam of light is focused laser light.

Claim 5. The method of Claim 1 further including a first step of treating the plant or animal tissue with at least one photo-active molecular agent, wherein the particular volume of the plant or animal tissue retains at least a portion of the at least one photo-active molecular agent.

Claim 6. The method of Claim 5 wherein the at least one photo-active molecular agent is selected from the group consisting of psoralen, 5-methoxypsoralen (5-MOP), 8-methoxypsoralen (8-MOP), 4,5',8-trimethylpsoralen (TMP), 4'-aminomethyl-4,5',8-trimethylpsoralen (AMT), 5-chloromethyl-8-methoxypsoralen (HMT), angelicin (isopsoralen), 5-methylangelicin (5-MIP), 3-carboxypsoralen, porphyrin, haematoporphyrin derivative (HPD), photofrin II, benzoporphyrin derivative (BPD), protoporphyrin IX (PpIX), dye haematoporphyrin ether (DHE), polyhaematoporphyrin esters (PHE), 13,17-N,N,N-dimethylethylethanolamine ester of protoporphyrin (PH1008), tetra(3-hydroxyphenyl)-porphyrin (3-THPP), tetraphenylporphyrin monosulfonate (TPPS1), tetraphenylporphyrin disulfonate (TPPS2a), dihaematoporphyrin ether, mesotetraphenylporphyrin, mesotetra(4N-methylpyridyl)porphyrin (T4MpyP), octa-(4-*tert*-butylphenyl)tetrapyrazinoporphyrazine (OTPP), phthalocyanine, tetra-(4-*tert*-butyl)phthalocyanine (t_4 -PcH₂), tetra-(4-*tert*-butyl)phthalocyanatomagnesium (t_4 -PcMg), chloroaluminum sulfonated phthalocyanine (CASPc), chloroaluminum phthalocyanine tetrasulfate (AlPcTS), mono-sulfonated aluminum phthalocyanine (AlSPc), di-sulfonated aluminum phthalocyanine (AlS2Pc), tri-sulfonated aluminum phthalocyanine (AlS3Pc), tetra-sulfonated aluminum phthalocyanine

20 (AlS4Pc), silicon phthalocyanine (SiPc IV), zinc II phthalocyanine (ZnPc), bis(di-isobutyl
octadecylsiloxy)silicon 2,3-naphthalocyanine (isoBOSINC), germanium IV
octabutoxyphthalocyanine (GePc), rhodamine 101 (Rh-101), rhodamine 110 (Rh-110), rhodamine
123 (Rh-123), rhodamine 19 (Rh-19), rhodamine 560 (Rh-560), rhodamine 575 (Rh-575),
rhodamine 590 (Rh-590), rhodamine 610 (Rh-610), rhodamine 640 (Rh-640), rhodamine 6G (Rh-
6G), rhodamine 700 (Rh-700), rhodamine 800 (Rh-800), rhodamine B (Rh-B), sulforhodamine
101, sulforhodamine 640, sulforhodamine B, coumarin 1, coumarin 2, coumarin 4, coumarin 6,
coumarin 6H, coumarin 7, coumarin 30, coumarin 47, coumarin 102, coumarin 106, coumarin
120, coumarin 151, coumarin 152, coumarin 152A, coumarin 153, coumarin 311, coumarin 307,
coumarin 314, coumarin 334, coumarin 337, coumarin 343, coumarin 440, coumarin 450,
coumarin 456, coumarin 460, coumarin 461, coumarin 466, coumarin 478, coumarin 480,
coumarin 481, coumarin 485, coumarin 490, coumarin 500, coumarin 503, coumarin 504,
coumarin 510, coumarin 515, coumarin 519, coumarin 521, coumarin 522, coumarin 523,
coumarin 535, coumarin 540, coumarin 540A, coumarin 548, 5-ethylamino-9-
diethylaminobenzo[a]phenoxazinium (EtNBA), 5-ethyl-amino-9-diethyl-
aminobenzo[a]phenothiazinium (EtNBS), 5-ethylamino-9-diethylaminobenzo[a]pheno-
selenazinium (EtNBSe), chlorpromazine, chlorpromazine derivatives, chlorophyll derivatives,
bacteriochlorophyll derivatives, metal-ligand complexes, tris(2,2'-bipyridine)ruthenium (II)
dichloride (RuBPY), tris(2,2'-bipyridine)rhodium (II) dichloride (RhBPY), tris(2,2'-
bipyridine)platinum (II) dichloride (PtBPY), pheophorbide a, merocyanine 540, vitamin D, 5-
amino-laevulinic acid, photosan, chlorin e6, chlorin e6 ethylenediamide, mono-L-aspartyl chlorin
e6, phenoxazine Nile blue derivatives, stilbene, stilbene derivatives, 4-(N-(2-hydroxyethyl)-N-

0002065 050509
methyl)-aminophenyl)-4'-(6-hydroxyhexylsulfonyl)stilbene (APSS), and standard biological dyes and stains.

Claim 7. The method of Claim 5 wherein the at least one photo-active molecular agent is at least one biogenic photo-active molecular agent that is specific to a particular tissue within the particular volume of plant or animal tissue.

Claim 8. The method of Claim 7 wherein the at least one biogenic photo-active molecular agent includes a segment selected from the group consisting of DNA, RNA, amino acids, proteins, antibodies, ligands, haptens, carbohydrate receptors or complexing agents, lipid receptors or complexing agents, protein receptors or complexing agents, chelators, and encapsulating vehicles.

Claim 9. The method of Claim 8 wherein the at least one biogenic photo-active molecular agent further includes a segment which is photo-activated when subject to light sufficient to promote a simultaneous two-photon excitation.

10
365050-2902200

Claim 10. The method of Claim 1 wherein the step of treating the particular volume of the plant or animal tissue with light sufficient to promote a simultaneous two-photon excitation of the at least one photo-active molecular agent contained in the particular volume of the plant or animal tissue includes the steps of:

- (a1) modulating light from a light source with a particular type of modulation, thereby producing a modulated light; and
 - (a2) treating the particular volume of the plant or animal tissue with the modulated light sufficient to promote a simultaneous two-photon excitation of the at least one photo-active molecular agent contained in the particular volume of the plant or animal tissue;
- and further including the steps of:
- (e) demodulating the detected energy signal with the particular type of modulation; and
 - (f) producing a demodulated energy signal which is characteristic of the particular volume of the plant or animal tissue.

Claim 11. The method of Claim 10 wherein the step of demodulating the detected energy signal with the particular type of modulation includes demodulating the detected energy signal at a frequency twice that of the particular type of modulation, thereby detecting the second harmonic of the particular type of modulation.

Claim 12. The method of Claim 10 wherein the demodulated energy signal which is characteristic of the particular volume of the plant or animal tissue represents a change in lifetime of at least one photo-activated molecular agent present in the particular volume of the plant or animal tissue.

Claim 13. A method for the imaging of a particular volume of material, wherein the material contains at least one photo-active molecular agent, the method comprising the steps of:

- (a) treating the particular volume of the material with light sufficient to promote a simultaneous two-photon excitation of at least one of the at least one photo-active molecular agent contained in the particular volume of the material;
- (b) photo-activating the at least one photo-active molecular agent in the particular volume of the material, thereby producing at least one photo-activated molecular agent, wherein the at least one photo-activated molecular agent emits energy;
- (c) detecting the energy emitted by the at least one photo-activated molecular agent; and
- (d) producing a detected energy signal which is characteristic of the particular volume of the material.

Claim 14. The method of Claim 13 wherein the material is selected from the group consisting of plant tissue and animal tissue.

Claim 15. The method of Claim 13 wherein the light sufficient to promote a simultaneous two-photon excitation of the at least one photo-active molecular agent is laser light.

Claim 16. The method of Claim 13 wherein the light sufficient to promote a simultaneous two-photon excitation of the at least one photo-active molecular agent is a focused beam of light.

Claim 17. The method of Claim 16 wherein the focused beam of light is focused laser light.

Claim 18. The method of Claim 13 further including a first step of treating the material with at least one photo-active molecular agent, wherein the particular volume of the material retains at least a portion of the at least one photo-active molecular agent.

Claim 19. The method of Claim 18 wherein the at least one photo-active molecular agent is selected from the group consisting of psoralen, 5-methoxypsoralen (5-MOP), 8-methoxypsoralen (8-MOP), 4,5',8-trimethylpsoralen (TMP), 4'-aminomethyl-4,5',8-trimethylpsoralen (AMT), 5-chloromethyl-8-methoxypsoralen (HMT), angelicin (isopsoralen), 5-methylangelicin (5-MIP), 3-carboxypsoralen, porphyrin, haematoporphyrin derivative (HPD), photofrin II, benzoporphyrin derivative (BPD), protoporphyrin IX (PpIX), dye haematoporphyrin ether (DHE), polyhaematoporphyrin esters (PHE), 13,17-N,N,N-dimethylethylethanolamine ester of protoporphyrin (PH1008), tetra(3-hydroxyphenyl)-porphyrin (3-THPP), tetraphenylporphyrin monosulfonate (TPPS1), tetraphenylporphyrin disulfonate (TPPS2a), dihaematoporphyrin ether, mesotetraphenylporphyrin, mesotetra(4N-methylpyridyl)porphyrin (T4MpyP), octa-(4-*tert*-

15

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

61

62

63

64

65

66

67

68

69

70

71

72

73

74

75

76

77

78

79

80

81

82

83

84

85

86

87

88

89

90

91

92

93

94

95

96

97

98

99

100

101

102

103

104

105

106

107

108

109

110

111

112

113

114

115

116

117

118

119

120

121

122

123

124

125

126

127

128

129

130

131

132

133

134

135

136

137

138

139

140

141

142

143

144

145

146

147

148

149

150

151

152

153

154

155

156

157

158

159

160

161

162

163

164

165

166

167

168

169

170

171

172

173

174

175

176

177

178

179

180

181

182

183

184

185

186

187

188

189

190

191

192

193

194

195

196

197

198

199

200

201

202

203

204

205

206

207

208

209

210

211

212

213

214

215

216

217

218

219

220

221

222

223

224

225

226

227

228

229

230

231

232

233

234

235

236

237

238

239

240

241

242

243

244

245

246

247

248

249

250

251

252

253

254

255

256

257

258

259

260

261

262

263

264

265

266

267

268

269

270

271

272

273

274

275

276

277

278

279

280

281

282

283

284

285

286

287

288

289

290

291

292

293

294

295

296

297

298

299

300

301

302

303

304

305

306

307

308

309

310

311

312

313

314

315

316

317

318

319

320

321

322

323

324

325

326

327

328

329

330

331

332

333

334

335

336

337

338

339

340

341

342

343

344

345

346

347

348

349

350

351

352

353

354

355

356

357

358

359

360

361

362

363

364

365

366

367

368

369

370

371

372

373

374

375

376

377

378

379

380

381

382

383

384

385

386

387

388

389

390

391

392

393

394

395

396

397

398

399

400

401

402

403

404

405

406

407

408

409

410

411

412

413

414

415

416

417

418

419

420

35 bacteriochlorophyll derivatives, metal-ligand complexes, tris(2,2'-bipyridine)ruthenium (II) dichloride (RuBPY), tris(2,2'-bipyridine)rhodium (II) dichloride (RhBPY), tris(2,2'-bipyridine)platinum (II) dichloride (PtBPY), pheophorbide a, merocyanine 540, vitamin D, 5-amino-laevulinic acid, photosan, chlorin e6, chlorin e6 ethylenediamide, mono-L-aspartyl chlorin e6, phenoxazine Nile blue derivatives, stilbene, stilbene derivatives, 4-(N-(2-hydroxyethyl)-N-methyl)-aminophenyl)-4'-(6-hydroxyhexylsulfonfyl)stilbene (APSS), and standard biological dyes and stains.

Claim 20. The method of Claim 18 wherein the at least one photo-active molecular agent is at least one biogenic photo-active molecular agent that is specific to a particular substance within the particular volume of material.

Claim 21. The method of Claim 20 wherein the at least one biogenic photo-active molecular agent includes a segment selected from the group consisting of DNA, RNA, amino acids, proteins, antibodies, ligands, haptens, carbohydrate receptors or complexing agents, lipid receptors or complexing agents, protein receptors or complexing agents, chelators, and encapsulating vehicles.

Claim 22. The method of Claim 21 wherein the at least one biogenic photo-active molecular agent further includes a segment which is photo-activated when subject to light sufficient to promote a simultaneous two-photon excitation.

Claim 23. The method of Claim 13 wherein the step of treating the particular volume of the material with light sufficient to promote a simultaneous two-photon excitation of the at least one photo-active molecular agent contained in the particular volume of the material includes the steps of:

(a1) modulating light from a light source with a particular type of modulation, thereby producing a modulated light; and

(a2) treating the particular volume of the material with the modulated light sufficient to promote a simultaneous two-photon excitation of the at least one photo-active molecular agent contained in the particular volume of the material;

and further including the steps of:

(e) demodulating the detected energy signal with the particular type of modulation; and

(f) producing a demodulated energy signal which is characteristic of the particular volume of the material.

Claim 24. The method of Claim 23 wherein the step of demodulating the detected energy signal with the particular type of modulation includes demodulating the detected energy signal at a frequency twice that of the particular type of modulation, thereby detecting the second harmonic of the particular type of modulation.

Claim 25. The method of Claim 23 wherein the demodulated energy signal which is characteristic of the particular volume of the material represents a change in lifetime of at least one photo-activated molecular agent present in the particular volume of the material.